

Contarctive properties of multifunctions related to uniformity

Pratip Chakraborty¹

University of Kaiserslautern

Kaiserslautern, DE 67663

chakrabo@mathematik.uni-kl.de

28/01/2005

1. INTRODUCTION

In this paper I will try to prove some contactive properties for multifunctions in uniformly complete metric spaces which are orbitally complete.

1.1. Basic definitions and notations. Let (X, U) be uniform space. A family $\{d_i : i \in I\}$ of pseudometrics on X with indexing set I , Is called an associated family for the uniformity U if the family

$$\beta = \{V(i, \varepsilon) : i \in I; \varepsilon > 0\}$$

where

$$v(i, \varepsilon) = \{(x, y) : x, y \in A, i \in I\}$$

is a sub-base for the uniformity U . We may assume that β itself is a base by adjoining finite intersection members of β , if necessary. The corresponding family of pseudometrics is called an associated family for U . An associated family for U will be denoted by p^* . Let A be a nonempty subset of a uniform space X . Define

$$D^*(A) = \sup\{d_i(x, y) : x, y \in A, i \in I\}$$

where

$$\{d_i : i \in I\} = p^*$$

Then D^* is called an augmented diameter of A . Furthermore A is said to be p^* -bounded if $D^*(A) < \infty$. Let

$$2^X = \{A : A \text{ is nonempty, closed and } p^*\text{-bounded subset of } X\}$$

For any non empty subsets of A and B of X , define

$$d_i(x, A) = \inf\{d_i(x, a) : a \in A\}, i \in I$$

$$H_i(A, B) = \max\{\sup d_i(a, B), \sup d_i(A, b)\} = \sup\{|d_i(x, A) - d_i(x, B)|\}$$

where $a \in A, b \in B$ and $x \in X$

It is well known that on 2^X , H_i is a pseudometric, called the induced Hausdorff pseudometric induced by $d_i, i \in I$.

Let (X, U) be a uniform space with an augmented associated family p^* . p^* also induces a uniformity U^* on 2^X defined by the base

$$\beta^* = \{V^*(i, \varepsilon) : i \in I, \varepsilon > 0\}$$

where

$$V^*(i, \varepsilon) = \{(A, B) : A, B \in 2^X, H_i(A, B) < \varepsilon\}.$$

The space $(2^X, U^*)$ is a uniform space called the hyperspace of (X, U) .

Let us review two following definitions.

Subfilter :

The collection of all filters on a given set x is denoted by $\Phi(X)$. An order relation

¹© My pleasure if you want to copy

is defined on $\Phi(X)$ by the rule $\mathfrak{F}_1 < \mathfrak{F}_2$ iff $\mathfrak{F}_1 \supset \mathfrak{F}_2$. If $\mathfrak{F}^* < \mathfrak{F}$, then $\mathfrak{F}^*$ is called a subfilter of $\mathfrak{F}$.

Orbital Completeness : Let (X, U) be a uniform space defined by $\{d_i : i \in I\} = p^*$. if $F : X \rightarrow 2^X$ is a multifunction, then

- (1) $x \in X$ is called a fixed point of F if $x \in Fx$; complete
- (2) an orbit of F at a point $x_0 \in X$ is a sequence $\{x_n\}$ given by

$$O(F, x_0) = \{x_n : x_n \in Fx_{n-1}, n = 1, 2, \dots\};$$

- (3) A uniform space X is called *F-orbitally complete* if every Cauchy filter which is a subfilter of an orbit of F at each $x \in X$ converges to a point of X .

Orbital Continuity : Let (X, U) be a uniform space and let $F : X \rightarrow X$ be a function. A single-valued function F is *orbitally continuous* if $\lim(T^{n_i}x) = u$ implies $\lim T(T^{n_i}x) = Tu$ for each $x \in X$.

2. CONTRACTION THEOREMS

Theorem 2.1. Let (x, U) be a F -orbitally complete Hausdorff uniform space defined by $\{d_i : i \in I\} = p^*$. and $(2^X, U^*)$ a hyperspace and let $F : X \rightarrow 2^X$ be a continuous function with Fx compact for each $x \in X$. Assume that

$$\min\{H_i(Fx, Fy)^r, d_i(x, Fx)d_i(y, Fy)^{r-1}, d_i(y, Fy)^r\}$$

$$+ a_i \min\{d_i(x, Fy), d_i(y, Fx)\} \leq [b_i d_i(x, Fx) + c_i d_i(x, y)]d_i(y, Fy)^{r-1}$$

for all $i \in I$ and $x, y \in X$, where $r \geq 1$ is an integer, a_i, b_i, c_i are real numbers such that $0 < b_i + c_i < 1$, then F has a fixed point.

proof : Let x_0 be an arbitrary point in X and consider the sequence defined by

$$x_1 \in Fx_0, x_2 \in Fx_1, \dots, x_n \in Fx_{n-1}, \dots$$

Let us assume that $d_i(x_n, Fx_n) > 0$ for each $i \in I$ and $n = 0, 1, 2, \dots$ (otherwise for some positive integer $n, x_n \in Fx_n$ as desired.)

Let $\mathcal{U} \in U$ be an arbitrary entourage. Since β is a base for U , there exists $V(i, \varepsilon) \in \beta$ such that $V(i, \varepsilon) \subseteq \mathcal{U}$. Now $y \rightarrow d_i(x_0, y)$ is continuous on the compact set Fx_0 and this implies that there exists $x_1 \in Fx_0$ such that $d_i(x_0, x_1) = d_i(x_0, Fx_0)$. Similarly, Fx_1 is compact so there exists $x_2 \in Fx_1$ such that $d_i(x_1, x_2) = d_i(x_1, Fx_1)$. Counting, we obtain a sequence $\{x_n\}$ such that $x_{n+1} \in Fx_n$ and $d_i(x_n, x_{n+1}) = d_i(x_n, Fx_n)$. For $x = x_{n-1}$, and $y = x_n$ by the condition of the theorem we have

$$\begin{aligned} & \min\{H_i(Fx_{n-1}, Fx_n)^r, d_i(x_{n-1}, Fx_{n-1})d_i(x_n, Fx_n)^{r-1}, d_i(x_n, Fx_n)^r\} \\ & + a_i \min\{d_i(x_{n-1}, Fx_n), d_i(y, Fx_{n-1})\} \leq [b_i d_i(x_{n-1}, Fx_{n-1}) \\ & + c_i d_i(x_{n-1}, x_n)]d_i(x_n, Fx_n)^{r-1} \end{aligned}$$

or since $d_i(x_n, x_{n+1}) = 0, x_n \in Fx_{n-1}$. Hence we have

$$\begin{aligned} & \min\{d_i(x_n, x_{n+1})^r, d_i(x_{n-1}, x_n)d_i(x_n, x_{n+1})^{r-1}\} \\ & \leq [b_i d_i(x_{n-1}, x_n) + c_i d_i(x_{n-1}, x_n)]d_i(x_n, x_{n+1})^{r-1} \end{aligned}$$

and it follows that

$$\begin{aligned} & \min\{d_i(x_n, x_{n+1})^r, d_i(x_{n-1}, x_n)d_i(x_n, x_{n+1})^{r-1}\} \\ & \leq (b_i + c_i)d_i(x_{n-1}, x_n)d_i(x_n, x_{n+1})^{r-1} \end{aligned}$$

is not possible (as $0 < b_i + c_i < 1$), we have

$$d_i(x_n, x_{n+1})^r \leq (b_i + c_i)d_i(x_{n-1}, x_n)d_i(x_n, x_{n+1})^{r-1}$$

or

$$d_i(x_n, x_{n+1})^r \leq k_i d_i(x_{n-1}, x_n)d_i(x_n, x_{n+1})^{r-1},$$

where $k_i = b_i + c_i, 0 < k_i < 1$.

proceeding in this mway we get

$$\begin{aligned} d_i(x_n, x_{n+1}) &\leq k_i d_i(x_{n-1}, x_n) \\ &\leq k_i^2 d_i(x_{n-2}, x_{n-1}) \\ &\vdots \\ &\leq k_i^n d_i(x_0, x_1) \end{aligned}$$

Hence we obtain

$$\begin{aligned} d_i(x_n, x_m) &\leq d_i(x_n, x_{n+1}) + d_i(x_{n+1}, x_{n+2}) + \dots + d_i(x_{m-1}, x_m) \\ &\leq (k_i^n + k_i^{n+1} + \dots + k_i^{m-1})d_i(x_0, x_1) \\ &\leq k_i^n (1 + k_i + \dots + k_i^{m-n-1}) \\ &\leq \frac{k_i^n}{1 - k_i} (1 + k_i + \dots + k_i^{m-n-1})d_i(x_0, x_1). \end{aligned}$$

Since

$$\lim_{n \rightarrow \infty} k_i^n = 0,$$

it follows that there exists $N(i, \varepsilon)$ such that $d_i(x_n, x_m) < \varepsilon$ and hence $(x_n, x_m) \in \mathcal{U}$ for all $n, m \geq N(i, \varepsilon)$. therefore the sequence $\{x_n\}$ is a Cauchy sequence in the d_i -uniformity on X .

let $S_p = \{x_n : n \geq p\}$ for all positive integer p and let β be the filter basis $\{S_p : p = 1, 2, \dots\}$. Then since $\{x_n\}$ is a d_i -cauchy sequence for each $i \in I$, it is easy to see that the filter basis β is a Cauchy filter in the uniform space (X, U) . To see this we first note that the family $\beta = \{V^*(i, \varepsilon) : i \in I, \varepsilon > 0\}$ is a base for U as $p^* = \{d_i : i \in I\}$. Now since $\{x_n\}$ is a d_i -Cauchy sequence in X , there exists a positive integer p such that $d_i(x_n, x_m) \leq \varepsilon$ for $m \geq p, n \geq p$. This implies that $S_p \times S_p \subseteq V^*(i, \varepsilon)$. Thus given any $\mathcal{U} \in U$, we can find an $S_p \in \beta$ such that $S_p \times S_p \subseteq \mathcal{U}$. Hence β is a Cauchy filter in (X, U) . Since (X, U) is a F -orbitally complete and Hausdorff space, $S_p \rightarrow z$ for some $z \in X$. consequently $F(S_p) \rightarrow Fz$ (follows from the continuity of F). Also

$$S_{p+1} \subseteq F(S_p) = \cup \{Fx_n : n \geq P\}$$

for $p = 1, 2, \dots$ it follows that $z \in Fz$. Hence z is a fixed point of F . This completes our task. $\square$

If we take $r = 1$ then the previous theorem becomes as the following theorem.

Theorem 2.2. *Let (x, U) be a F -orbitally complete Hausdorff uniform space defined by $\{d_i : i \in I\} = p^*$. and $(2^X, U^*)$ a hyperspace and let $F : X \rightarrow 2^X$ be a continuous function with Fx compact for each $x \in X$. Assume that*

$$\begin{aligned} &\min\{H_i(Fx, Fy), d_i(x, Fx)d_i(y, Fy), d_i(y, Fy)\} \\ &+ a_i \min\{d_i(x, Fy), d_i(y, Fx)\} \leq b_i d_i(x, Fx) + c_i d_i(x, y) \end{aligned}$$

for all $i \in I$ and $x, y \in X$, where a_i, b_i, c_i are real numbers such that $0 < b_i + c_i < 1$, then F has a fixed point.

We denote that if F is a single valued function on X , then we can write $d_i(Fx, Fy) = H_i(Fx, Fy), x, y \in X, i \in I$. Thus we obtain the following theorem as a consequence of the second theorem where we assume $r = 1$.

Theorem 2.3. *Let (x, U) be a T-orbitally complete Hausdorff uniform space and let $T : X \rightarrow X$ be a T-orbitally continuous function satisfying*

$$\begin{aligned} & \min\{d_i(Tx, Ty), d_i(x, Tx)d_i(y, Ty), d_i(y, Ty)\} \\ & + a_i \min\{d_i(x, Ty), d_i(y, Tx)\} \leq b_i d_i(x, Tx) + c_i d_i(x, y) \end{aligned}$$

for all $i \in I$ and $x, y \in X$, where a_i, b_i, c_i are real numbers such that $0 < b_i + c_i < 1$, then T has a unique fixed point whenever $a_i > c_i > 0$.

proof Define a function F of X into 2^X by putting $FX = \{Tx\}$ for all $x \in X$. it follows that F satisfies the conditions of second theorem. Hence T has a fixed point.

Now if $a_i > c_i > 0$, we show that T has a unique fixed point. Assume that T has two fixed points Z and w which are distinct. Since $d_i(z, Tz) = 0$ and $d_i(w, Tw) = 0$, then by the condition of the previous theorem,

$$a_i \min\{d_i(z, Tw), d_i(w, Tz)\} \leq c_i d_i(z, w)$$

or

$$a_i d_i(z, w) \leq c_i d_i(z, w),$$

$$d_i(z, w) \leq \frac{c_i}{a_i} d_i(z, w)$$

which is impossible. Thus if $a_i > c_i > 0$, then T has a unique fixed point in X . And that is what we wanted. $\square$

COROLLARY

Let T be an orbitally continuous self-map of a T-orbitally complete uniform space (X, U) satisfying the condition

$$\begin{aligned} & \min\{d_i(Tx, Ty), d_i(x, Tx), d_i(y, Ty)\} \\ & - \min\{d_i(x, Ty), d_i(y, Tx)\} \leq b_i d_i(x, Tx) + c_i d_i(x, y), \end{aligned}$$

$x, y \in X, i \in I$ and $0 < b_i + c_i < 1$. Then for each $x \in X$, the sequence $\{T^n x\}$ converges to a fixed point of T .

I want to remark that this theorem came to my mind as I thought that the result obtained by Dhage[2] can be extended with more generalisation from metric space to spaces of arbitrary uniform structure. It is obvious that Theorem 1 and Corollary 1 of his paper will follow from our last theorem and the corollary if we replace the uniform space (X, U) by a metric space.

REFERENCES

- [1] Acharya S P, Some results on fixed point in uniform space, *Yokohama Math. J.* 23(1) (1974) 105-116
- [2] Dhage B C, Some results for the map with a nonunique fixed point, *Indian J. Pure Appl. Math.* 16(3)(1985) 245-256
- [3] Kelly J L, *General Topology* (Van Nostrand Company Inc) (1955)
- [4] Thron W J, *Topological structures* (New York: Holt, Rinehart and Wintson) (1966)